

BMJ Open Infant formula supplemented with milk fat globule membrane compared with standard infant formula for the cognitive development of healthy term-born formula-fed infants: protocol for a randomised controlled trial

Jacqueline F Gould ^{1,2}, Robert A Gibson,^{1,3,4} Lisa N Yelland ^{1,5}, John Colombo,⁶ Andrew J McPhee,^{1,7} Sophie Gallier,^{8,9} Rachel M Roberts,² D. Jill Shaddy,¹⁰ Jana Bednarz,¹¹ Maria Makrides¹²

To cite: Gould JF, Gibson RA, Yelland LN, *et al.* Infant formula supplemented with milk fat globule membrane compared with standard infant formula for the cognitive development of healthy term-born formula-fed infants: protocol for a randomised controlled trial. *BMJ Open* 2024;**14**:e083399. doi:10.1136/bmjopen-2023-083399

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjopen-2023-083399>).

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2023-083399>).

Received 19 December 2023
Accepted 13 June 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jacqueline F Gould;
jacqueline.gould@sahmri.com

ABSTRACT

Introduction Milk fat globule membrane (MFGM) is a complex lipid–protein structure in mammalian milk and human milk that is largely absent from breastmilk substitutes. The objective of this trial is to investigate whether providing infant formula enriched with MFGM versus standard infant formula improves cognitive development at 12 months of age in exclusively formula-fed full-term infants.

Methods and analysis This is a randomised, controlled, clinician-blinded, researcher-blinded and participant-blinded trial of two parallel formula-fed groups and a breastfed reference group that were recruited in the suburban Adelaide (Australia) community by a single study centre (a medical research institute). Healthy, exclusively formula-fed, singleton, term-born infants under 8 weeks of age were randomised to either an MFGM-supplemented formula (intervention) or standard infant formula (control) from enrolment until 12 months of age. The reference group was not provided with formula. The primary outcome is the Cognitive Scale of the Bayley Scales of Infant Development, Fourth Edition (Bayley-IV) at 12 months. Secondary outcomes are the Bayley-IV Cognitive Scale at 24 months, other Bayley-IV domains (language, motor, emotional and behavioural development) at 12 and 24 months of age, infant attention at 4 and 9 months of age, parent-rated language at 12 and 24 months of age, parent-rated development at 6 and 18 months of age as well as growth, tolerance and safety of the study formula. To ensure at least 80% power to detect a 5-point difference in the mean Bayley-IV cognitive score, >200 infants were recruited in each group.

Ethics and dissemination The Women's and Children Health Network Human Research Ethics Committee reviewed and approved the study (HREC/19/WCHN/140). Caregivers gave written informed consent prior to enrolling in the trial. Findings of this study will be disseminated through peer-reviewed publications and conference presentations.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is a robust randomised controlled trial of infant formula enriched with milk fat globule membrane for exclusively formula-fed infants.
- ⇒ Longitudinal assessments across early development are comprehensive.
- ⇒ Methods were adapted to coronavirus-related restrictions prior to trial commencement.
- ⇒ Loss to follow-up after enrolment into the trial may contribute to risk of attrition bias.

Trial registration number ACTRN12620000552987; Australian and New Zealand Clinical Trial Registry: anzctr.org.au.

INTRODUCTION

Exclusive breast milk is considered the ideal source of nutrition for infants up to 6 months of age with suggested benefits to general health,¹ allergies,² infection,² growth and body composition,² for infants who are breastfed compared with infants who are formula-fed. In particular, breastmilk appears to confer benefits to general neurodevelopment,^{1 3} behaviour,³ and cognitive abilities, as captured by intelligence tests.^{2–5}

Breast milk contains a range of important nutrients for health and neurodevelopment^{6–8} that differs to formula and may contribute to the cognitive and other benefits seen in breastfed infants.^{1 2 4 5} Recent interest has been drawn to the milk fat globule membrane (MFGM), a structurally and compositionally complex membrane of bioactive compounds. MFGM is naturally occurring as the membrane of the fat globule

in breast milk but is only present in trace amounts in most infant formula where the fat source is mainly vegetable oils.⁹ It has recently become possible to produce MFGM of sufficient quality and quantity to be included in infant formulas, and multiple trials have tested the effects of giving infants MFGM.^{10–17 18–21} The results from these studies indicated that supplementation of bovine milk-derived MFGM was safe and may benefit cognition, however, results were mixed partially due to heterogeneity of intervention (timing, duration and inclusion of other components in combination with MFGM), outcomes and study quality (in relation to limitations such as small samples or MFGM intake of the control group).^{10–21} Most noteworthy is that technician-administered tests have suggested benefits to cognitive development at 12¹⁰ months of age in the most robust trial of 160 formula-fed term-born infants, although by 6 years of age, there were no group differences.²² Additionally, this benefit to cognitive development was not detected in other trials with assessments at 2, 3, 4, 6, 12 or 18 months of age.^{11–13}

We report the design of a new, large, robust trial with the aim to determine the effect of MFGM-supplemented formula compared with standard formula for exclusively formula-fed full-term infants from <2 months of age until 12 months of age on their cognitive development. We hypothesise that supplementing the formula of non-breastfed infants with MFGM will improve cognitive scores at 12 months of age compared with infants receiving standard infant formula.

METHODS AND ANALYSIS

Study design

The Infant nutrition with Milk fAt Globule membrane for Infant cognition in Early life (IMAGINE) trial is a double-blind randomised controlled trial of infants that are not receiving any breastmilk, and a non-randomised breastfed reference group. The IMAGINE Trial Coordinating Centre was the South Australian Health and Medical Research Institute (SAHMRI) Women and Kids theme, based at the Women's and Children's Hospital (Adelaide, South Australia). A purpose-built web-based (Research Electronic Data Capture; REDCap) management system is used to collect and store data.

Trial status

The IMAGINE trial recruitment began on 14 May 2020. As of November 2023, randomised and enrolment is complete with a total of 620 participants. Completion of participant follow-up assessments to 24 months of age is expected by 30 June 2024.

Participants and sample selection

Infants up to 2 months (60 days) of age were eligible to enrol.

Inclusion criteria

- ▶ Singleton birth.

- ▶ Full-term birth (37⁺⁰ to <41⁺⁶ weeks).
- ▶ Appropriate birth weight (>5th and <95th percentile: boys ≥2604 and ≤4215 g, girls ≥2532 and ≤4041 g).²³
- ▶ The legal caregiver was able to provide written informed consent.
- ▶ The caregiver agreed not to enrol the infant in another interventional study that may affect infant growth or development.
- ▶ The family lived within metropolitan Adelaide or was willing to travel to appointments.
- ▶ English was the primary language spoken at home.
- ▶ Formula-fed groups only
 - Exclusively formula fed at the time of screening (anytime up to 60 days of age), where exclusively formula fed was defined as no breastmilk for at least 24 hours prior to consent, and caregivers had no plans to retry breastfeeding.
 - No medically diagnosed allergy or intolerance to lactose, soy, fish or cow's milk protein.
- ▶ Breastfed reference group only
 - Infants were at least 52 days of age at enrolment
 - Infants were exclusively fed breastmilk at enrolment (52–60 days of age).
 - Caregivers planned to feed breastmilk until the infant was at least 12 months of age.

Exclusion criteria

- ▶ Severe congenital or metabolic disease, or congenital malformation, major birth defect or any other condition likely to interfere with
 - the ability to ingest food.
 - normal growth and development.
 - evaluation of the infant
- ▶ Hypoxic ischaemic encephalopathy.
- ▶ Birth mother had pre-existing type 1 diabetes.
- ▶ Known substance or alcohol misuse during pregnancy.
- ▶ Infant was immunocompromised (according to a doctor's diagnosis of immunodeficiency such as combined immunodeficiency's, DiGeorge syndrome, Wiskott-Aldrich syndrome, severe congenital neutropenia and secondary immunodeficiencies linked to HIV infection, Down syndrome or others).
- ▶ Already participating in an intervention study that may influence development.

Screening and enrolment

There was no advertising for the IMAGINE trial. Potentially eligible families were not informed about a formula study unless they had already confirmed their infant was exclusively formula-fed, or exclusively fed breastmilk. Families with potentially eligible infants were screened for the study through a state-wide community nurse service that performs routine health checks for all infants between 1 and 4 weeks of age in South Australia, and other SAHMRI research activities where infants feeding practices were already being collected (see [figure 1](#)). Families with singleton term-born infants that indicated exclusive formula feeding at the health check were




Figure 1 Participant flow from screening to enrolment and randomisation in the trial. SAHMRI, South Australian Health and Medical Research Institute.

subsequently called to confirm feeding practices, and if still exclusively formula feeding, they were asked if they would like to receive information about a SAHMRI study. Some SAHMRI Women and Kids studies collected infant feeding practices²⁴ and allowed screening for term-born singleton infants under 2 months of age that were either exclusively formula-fed (formula-fed groups) or exclusively fed breastmilk (reference group).

SAHMRI staff contacted potentially eligible families to determine plans for ongoing feeding practices (breastmilk and/or infant formula), complete further screening and then (if eligibility criteria were met) explain study details and send an information sheet for review (see the online supplemental file 1 for the formula-fed group and breastfed reference group Participant Information and Consent Forms). Caregivers were given time to read the study information and discuss the study with family members and healthcare providers before being contacted by SAHMRI staff to potentially book a virtual enrolment appointment. A legal caregiver provided written informed e-consent through Adobe Sign or the REDCap e-consent module, both of which could be used on a basic smartphone,²⁵ prior to enrolling. Infants remained in the study until they completed the 24-month assessment (study exit).

Randomisation and blinding

On enrolment, infants were allocated a unique participant identification code (study ID) in the REDCap management system. Formula-fed infants were randomised to one of four colour-coded formula groups using a computer-generated randomisation schedule using a balanced variable block design. Infant formula was provided to participants in cans with plain packaging that were identical with the exception that the colours of the labels differed (for the four coloured groups). Two of the colour groups received the intervention formula, and two of the colour groups received the control formula (total of four colour groups to ensure blinding). Randomisation was stratified by infant sex, and infant age at enrolment (<28 days or ≥28 days) with balanced variable blocks of size 4 and 8. Both participants, their caregivers and all study staff were blinded to randomised allocation. The randomisation list was held by an independent statistician who could see which colour formula the infant was assigned to. Two custodians that did not have any contact with the study participants or have access to the data collected for the study held the randomisation list, with the unblinded group allocation. In the event of a medical or other emergency in which knowledge of group allocation was necessary, the custodians were available to provide emergency unblinding to the treating physician and the family.

The reference group of breastfed infants was not randomised or blinded to being the reference group.

Study intervention

Randomised infants received study formula manufactured and provided by Fonterra Co-Operative Group

Limited (New Zealand) from within a week of enrolment (≤2 months of age) until 12 months of age. The study formulas were cow's milk based and were matched for energy, fats, proteins, carbohydrates and key nutrients, except for MFGM (see table 1). The intervention formula contained a minimum ganglioside concentration of 17.9 mg/100 g, manufactured using bovine MFGM-rich ingredient as a source of gangliosides and milk phospholipids (MFGM Lipid 100; NZMP, Fonterra) from anhydrous milk fat production. The control formula was manufactured with the same macro and micronutrient composition but without the MFGM-enriched ingredient. Intervention and control study formulas were in powdered form to be reconstituted to deliver 66 kcal per 100 mL.

Study formulas were delivered to participants' residences packed in 900 g cans labelled with clear instructions for preparing the formula. Frequency and dose of study formula feeding were ad libitum at the discretion of the caregiver. Caregivers could request regular deliveries of formula as needed and study staff enquired about formula delivery requirements at each study contact before 12 months. Caregivers were instructed to feed study formula as the sole source of infant formula through to 12 months of age, unless otherwise instructed by their physician. As per the Australian National Health and Medical Research Council guidelines for infant feeding,²⁶ caregivers of infants in the breastfed reference group were encouraged to feed breastmilk as the sole source of milk until 6 months of age and to continue breastfeeding until the infant was 12 months. Reference group families were offered appointments with a lactation consultant as needed prior to the infants reaching 6 months of age but were not provided with study product, regardless of timing of breastfeeding cessation. As per Australian guidelines, caregivers of all infants (randomised and reference groups) were advised to introduce complementary feeding around 6 months of age,²⁶ unless otherwise instructed by their physician.

Outcomes and measures

At enrolment, caregivers underwent a virtual appointment, and then received phone calls 4, 7, 14 and 21 days after enrolment, completed surveys and were invited to attend study visits (see figure 2 for ages and timelines of study contact points). Families were invited to attend face-to-face appointments when the infant was 4, 9, 12 and 24 months of age. As the face-to-face assessment at 4 and 9 months of age was conducted to collect an exploratory, secondary outcome, appointments were conducted virtually (without the face-to-face component of the assessment) if caregivers preferred. Online questionnaires were emailed or texted (dependent on caregiver preference) to families when the infant turned 3, 5, 6, 12, 18 and 24 months of age. See online supplemental table for a summary of study milestones and information collected. Infants in the formula-fed groups and the reference group underwent all assessments, with the exception that formula-fed groups were not asked about breastfeeding,

Table 1 Nutrient composition of study formulae

		Intervention formula		Control formula	
		Per 100 g	Per 100 mL	Per 100 g	Per 100 mL
Energy	kJ	2091.0	276.0	2091.0	276.0
Energy**	cal	500	66.0	500	66.0
Protein	g	10.9	1.4	10.9	1.4
Fat, total	g	26.52	3.5	26.52	3.5
Linoleic acid	mg	4807	635	4807	635
α -linolenic acid	mg	427	56	427	56
Docosahexaenoic acid (DHA)	mg	60	8	60	8
Arachidonic acid (ARA)	mg	76	10	76	10
Carbohydrate	g	54.4	7.2	54.4	7.2
Calcium	mg	370	49	370	49
Phosphorus	mg	210	28	210	28
Sodium	mg	142	19	142	19
Potassium	mg	560	74	560	74
Chloride	mg	350	46	350	46
Magnesium	mg	50	6.6	50	6.6
Iron	mg	6	0.8	6	0.8
Zinc	mg	4.6	0.61	4.6	0.61
Copper	μ g	410	54	410	54
Manganese	μ g	80	11	80	11
Iodine	μ g	90	12	90	12
Selenium	μ g	15	2	15	2
Vitamin A	μ g-RE	500	66	500	66
Vitamin D3	μ g	7.1	0.94	7.1	0.94
Vitamin E (α -Tocopherol acetate)	mg α -TE	9	1.2	9	1.2
Vitamin K1	μ g	50	6.6	50	6.6
Thiamin (vitamin B1)	μ g	500	66	500	66
Riboflavin (vitamin B2)	μ g	1000	132	1000	132
Niacin (vitamin B3)	μ g	4000	528	4000	528
Pantothenic acid (vitamin B5)	μ g	3000	396	3000	396
Vitamin B6	μ g	350	46	350	46
Biotin	μ g	15	2	15	2
Folic acid	μ g	85	11	85	11
Vitamin B12	μ g	1.7	0.23	1.7	0.23
Vitamin C	mg	81.2	10.7	81.2	10.7
Choline	mg	100	13	100	13
Taurine	mg	40	5.3	40	5.3
L-carnitine	mg	10	1.3	10	1.3
Inositol	mg	51.7	6.8	51.7	6.8
Nucleotides	mg	19.7	2.6	19.7	2.6
Galacto-oligosaccharide (GOS)	g	4.1	0.54	4.1	0.54




Figure 2 Age of participants at study milestones. ¹Timing of phone calls are based on days since enrolment, therefore infant age at phone calls varied. ²Optional face-to-face visit or virtual appointment. ³Primary outcome collected at 12-month appointment.

and the reference group was not asked about use of study formula.

Primary outcome: cognition at 12 months

The primary outcome is infant cognition at 12 months of age, as assessed with the Cognitive Scale score of the Bayley Scales of Infant and Toddler Development, Fourth Edition, Australian and New Zealand (Bayley-IV). The Cognitive Scale score evaluates sensorimotor

development, exploration, manipulation, object relatedness, concept formation, memory and simple problem-solving.²⁷ The Cognitive Scale score was age standardised to a mean of 100 and SD of 15 and can be considered an early indicator of IQ. Scores range from 45 to 155 where higher scores reflect more advanced performance, and development can be classified as normal (85–115), impaired (<85) or accelerated (>115). The Cognitive

Scale is administered at a face-to-face appointment by a psychologist, trained health professional or trained technician. Appointments were conducted at clinic rooms at the Women's and Children's Hospital, SAHMRI or the family home, depending on the preference of the family.

Secondary outcomes assessed by study staff

Other Bayley-IV Subscales

The Bayley-IV includes assessor-administered Language and Motor Scales Scores, in addition to the Cognitive Scale.²⁷ The Language Scale is a composite of receptive communication (verbal comprehension, vocabulary) and expressive communication (babbling, gesturing and utterances) abilities. The Motor Scale evaluates both gross (big, large movements such as rolling over or walking) and fine (small movements and manipulations using small muscles, such as the fingers or lips) motor functioning. All scales are administered when children reached 12 and 24 months of age and scores are age standardised.

Attention

The attention assessments required a face-to-face appointment at the hospital clinic room due to the equipment needed. The assessments were an exploratory outcome (to be published separately from the primary results paper) and, therefore, optional for family's willing and able to attend, ideally within ± 3 days of the infant turning 4 months of age and within ± 7 days of the infant reaching 9 months of age. Assessments were conducted in a plain quiet clinic room according to a standard operating procedure.^{28–30} Infants were seated on a caregiver's lap or in a highchair in front of a computer monitor while their face was video recorded and heart rate was monitored.

The first task assessed visual habituation to repeatedly presented stimuli. Habituation is a longstanding and widely accepted measure of non-associative learning, and the visual habituation task assessed the relative speed with which infants encoded a visual stimulus. Infants were presented with a single stimulus (a woman's face^{30 31} over multiple trials interspersed with a blank screen while their attention to the stimulus was coded live).²⁸ Infant fixation (looking) declines with repeated exposure due to information processing or encoding (ie, learning). Once looking had declined to a predetermined threshold, the infant was considered 'habituated' to the stimulus. Subsequently, the infant was presented with the habituated stimulus and a novel stimulus (another woman's face³⁰ simultaneously paired). Infants who had successfully encoded the initial (ie, habituated) stimulus look preferentially at the novel stimulus. Along with assessing the learning that has occurred during habituation, the preference for the novel stimulus serves as an indication of recognition memory. The heart rate recording was synchronised with presentation of the stimulus and the video recording of infant looks to and from the stimulus to parse specific heart rate-defined phases of attention; orienting (initial look at the stimulus, prior to stable

deceleration of the heart rate), sustained attention (heart rate deceleration, reflecting encoding or processing of the stimulus) and attention termination (looking at the stimulus but disengaging, heart rate deceleration has ended).^{28 32–34} Outcomes for comparison were fixation duration, novelty preference, heart rate and proportion of looking spent in the different phases of attention.²⁸

The second task assessed ability to disengage attention or visual reaction time.^{35–37} The infant was presented with a stimulus (small colourful shape) in the middle of the monitor until their attention had been drawn to the centre of the screen. Another similar stimulus is then presented in the periphery of the infant's visual field either after the withdrawal of centre stimulus (ie, there was a 'gap' between the central stimulus' disappearance and the presentation of the peripheral stimulus) or prior to the withdrawal of the centre stimulus (ie, the two stimuli 'overlap'). The main outcome for comparison was the latency for the infant to initiate a correct shift in gaze from the initial stimulus location to the peripheral stimulus in the gap and overlap conditions.

Growth

Length (or height), weight and head circumference are assessed at 4, 9, 12 and 24 months of age. Where study staff are unable to conduct assessments, measurements taken by healthcare professionals (from routine check-ups or vaccination clinics) are obtained. Measurements will be converted to Z (SD) scores appropriate for age and sex.³⁸ Growth measurements at each time point as well as trajectory of growth across the study will be compared between the groups.

Secondary outcomes reported by parent-completed questionnaires or interview

Bayley-IV

The Bayley-IV Social-Emotional and Adaptive Behaviour Scales are administered when the infant is 12 and 24 months of age. The Social-Emotional Scale assesses milestones such as engagement and use of a range of emotions, experiences and expressions. The Adaptive Behaviour Scale captures self-direction abilities such as learning to feed oneself, crawling, walking and toileting, and communicating for basic needs.²⁷

Ages and Stages Questionnaire, Third Edition

The Ages and Stages Questionnaire, Third Edition (ASQ-3) ASQ-3 is a screen of early developmental milestones in five domains: communication skills, gross motor skills, fine motor skills, problem-solving skills and personal-social development.³⁹ The ASQ-3 was administered as infants turned 6 and 18 months of age.

MacArthur-Bates Communicative Development Inventory, Third Edition

The MacArthur-Bates Communicative Development Inventory, Third Edition (MCDI-3) is a well-validated and standardised parent-reported measure of early verbal and non-verbal communication.⁴⁰ The Words and

Gestures form was administered at 12 months to capture words understood and spoken as well as gestures used to communicate. The Words and Sentences form is used at 24 months of age to assess words produced and how they were used, early grammar and words understood. The MCDI-3 includes an online scoring programme with age-standardised norms. We modified some words for use in an Australian sample (eg, 'coin' rather than 'penny' and 'mummy' rather than 'mommy').⁴⁰

Feeding information and compliance

Feeding practices were collected at each call, survey and visit up to and including the visit at 12 months of age to determine compliance and identify group imbalances. Ongoing use of study formula, non-study formula, introduction of complementary feeds and use of dietary supplements were collected. Caregivers were asked how many bottles of study formula their infant had been provided in the previous 3 days, how much made-up formula was generally in each bottle and how much of the bottle was typically discarded (left over-after feeding). Reasons for and number of days of providing non-study formula (to the formula-fed or breastfed groups) during the intervention period were asked, as was the date and reason for ceasing study formula or breastfeeding. If non-study formula was provided to an infant from the formula-fed groups, or if any formula was fed to infants in the breastfed group, the formula was recorded as being either cows' milk based, soy based or hypoallergenic.

Infants in both the formula-fed groups and breastfed reference group were considered compliant if the number of non-compliant days (days where one or more feeds are non-study formula, respectively) was <10% between study commencement (7 days after enrolment) and 6 months of age.

Illness and serious adverse events

At each visit and survey, caregivers are asked to list consultations (visits or phone calls) with a healthcare professional (eg, a doctor, community nurse, hospital or parent helpline), symptoms that prompted the consultation, and any medications prescribed as well as any fevers (not in relation to a vaccination). At 24 months, caregivers provide neurological or chronic medical conditions the infant had been diagnosed with (for example, cerebral palsy or allergies). Potential serious adverse events (SAEs) are confirmed through review of infant medical records, and were defined as any one of the following

- ▶ Requirement of inpatient hospitalisation for 6 or more hours or prolongation of existing hospitalisation.
- ▶ Persistent or significant disability/incapacity.
- ▶ Life-threatening condition.
- ▶ Death.

Tolerance

At each visit and survey during the intervention, caregivers were asked about spilling and vomiting after feeding, fussiness and crying, faecal characteristics (rated

with a Bristol Stool Chart,⁴¹ and sleeping patterns to indicate tolerance). Formula-fed group caregivers were also asked about use of thickeners or other additives when preparing the study formula.

Background information and characteristics

Characteristics were collected to describe the sample, identify possible randomisation group imbalances and for inclusion of prespecified covariates in analysis models. At enrolment, a range of parental and infant sociodemographic and health information was collected through scripted interview. Where possible, caregivers provided their infants' health and development booklet (a hard copy booklet given to infants for health professionals to record health information) for verification. Recorded maternal information included age, country of birth, cultural ethnicity, education, working status in the previous 12 months, height, weight and prepregnancy weight. If the primary carer was not the biological mother, we collected information about education and working status in the last 12 months. Pregnancy and neonatal characteristics collected included prenatal smoking or alcohol use, mode of delivery, parity, anthropometrics at birth, neonatal illness or medical conditions, type of first feed, feeds while in hospital, any exclusive breastfeeding, alcohol use while breastfeeding and use of infant dietary supplements or thickeners prior to enrolment.

Stimulation within the home environment has an important role in the cognitive, social and emotional developmental outcomes of infants and is measured with a revised version of the Home Screening Questionnaire⁴² at 12 and 24 months of age. Childcare and number of adults and children living with the infant may be considered when comparing neurodevelopment outcomes or communicable illnesses.

Sample size

A sample size of 143 infants per formula-fed group (total 286) provides 80% power (two-tailed alpha 0.05) to detect a 5-point mean difference in the primary outcome of 12-month Bayley-IV Cognitive Scale between the randomised formula-fed groups. A 5-point difference was chosen as the minimum clinically important difference as it is similar in magnitude to the effect sizes that led to iron fortification of infant cereals,⁴³ and the removal of lead from petrol and the environment.⁴⁴ The sample size calculation is based on the Bayley-IV scales having a mean of 100 and SD of 15 in the general population at any time point. In the term-born, healthy population eligible for this study, the SD is likely to be lower (closer to 13,⁴⁵ which would mean a sample size of 144 per formula-fed group confers 90% power to detect a 5-point mean difference in the Cognitive Scale). A sample size target of 200 infants randomised per group allows up to 28% loss to follow-up.

As a secondary analysis, there will be a three-way comparison between the two randomised formula-fed groups and the breastfed reference group. Pairwise comparisons involving the reference group are powered

to detect a similar difference (90% power for 5 points on the Bayley-IV Cognitive Scale, SD of 13, two-tailed $\alpha=0.05$ for each comparison) to the primary outcome comparison.

Data management and analysis plan

Data are entered into REDCap and hosted on SAHMR's secure servers. REDCap uses a MySQL database via a secure web interface with data checks during data entry to ensure quality. REDCap includes a complete suite of features to support compliance with the Health Insurance Portability and Accountability Act of 1996, including a full audit trail, user-based privileges and integration with the institutional LDAP server.

All analyses will be conducted according to a prespecified statistical analysis plan⁴⁶ approved by the IMAGINE Steering Committee. Analyses will not commence until completion of data collection and blinded data cleaning, and the database is locked. All participants will be analysed according to the group into which they were randomised (intention-to-treat principle) in the primary analyses, regardless of compliance. Blinded treatment group codes (eg, group A or B) will be made available to the trial statistician and analyses will be performed using these group codes to allow analyses to be performed blinded to treatment group. Blinding will be broken after the results have been reviewed by the investigators.

Differences between the two randomised groups on outcomes measured at a single time point will be assessed using linear regression models for continuous outcomes, and log binomial regression models for binary outcomes and negative binomial regression for count outcomes. Differences between the two randomised groups on outcomes measured over time will be assessed using linear mixed effects models for continuous outcomes and log binomial models with generalised estimating equations for binary outcomes. The models will include terms for randomised group, time (treated as categorical due to measurements being collected at a small number of specific timepoints) and the interaction between group and time, with adjustment for randomisation strata (infant sex and infant age at enrolment) in adjusted analyses. Both adjusted and unadjusted analyses will be performed, with the adjusted results used to draw conclusions about the effect of the intervention on the outcomes of interest. Results will be presented as differences in means for continuous outcomes, or relative risks for binary outcomes, or incidence rate ratios for count outcomes at each time point where applicable, with 95% CIs and two-sided *p* values. Statistical significance will be assessed at the 5% level and no adjustment will be made for the number of analyses planned, as a single primary outcome has been pre-specified for the study. No interim analyses are planned.

Secondary per-protocol analyses will be conducted for the primary outcome and secondary outcomes from the Bayley-IV assessments and will include all participants consented into the study with no major protocol violations

and adequate compliance (<10% non-compliant days, with additional secondary analyses performed for <15% and <25% non-compliant days).

Comparisons involving the breastfed reference group will be based on the same statistical models as described above, but with additional adjustment for prespecified potential confounders. A global test will be performed for the effect of group (intervention, control or reference) followed by post hoc pairwise comparisons with no adjustment for multiple comparisons due to the exploratory nature of these analyses. Only pairwise comparisons involving the reference group (ie, intervention vs reference group, and control vs reference group) will be reported from these models.

Data collected on participants up to the point of withdrawal or loss to follow-up will be included in the analysis. Where data are missing, multiple imputation will be used to create 100 complete datasets for analysis of the primary and secondary Bayley-IV related outcomes using the fully conditional specification method, performed separately by treatment group.⁴⁷ Imputed datasets will include all surviving infants. Infants who were missing scores on psychological assessments because they were unable to complete the assessment for neurological reasons will be reviewed by the Chief Investigator and the Clinical Psychologist (blinded to treatment group) to determine whether assigning the lowest possible score is appropriate. All analyses will be performed blinded on both the raw and imputed data, with conclusions to be drawn based on the results using the imputed data.

Data monitoring and safety

The Steering Committee met monthly to monitor participants screened, enrolled/randomised, visits due/overdue/missed, adverse and SAEs), tolerance and compliance, product inventory and participant communication logs. An external monitor reviewed and verified all study documents and procedures as well as all signed consent forms, and a random selection of full participant records according to a predefined monitoring plan. A neonatologist reviewed all SAEs as well as monthly reports of illnesses and tolerance throughout the trial.

A Data Safety and Monitoring Committee comprising a paediatrician, a clinical researcher, and a biostatistician that were not otherwise involved in the study followed a formal charter to safeguard the interests of participants. The committee met quarterly to review adverse event data, SAEs as well as the indicators of dietary tolerance, growth and general well-being of the study groups to determine likelihood that involvement in the trial could have contributed to an event, and recommend whether or not to continue the trial.

Patient and public involvement

Neither patients nor the public were directly involved in the development of the research question or design of this study. Prior to the commencement of study recruitment, there was a COVID-19 outbreak, that necessitated

reducing face-to-face contact to minimise opportunities to spread the virus. A Community Board advisory group, comprising parents as well as clinicians and researchers specialising in paediatrics, was consulted on the revision of planned study procedures to safely conduct research activities during the COVID-19 outbreak.²⁵

Ethics and dissemination

The IMAGINE trial is being conducted in accordance with the Australian National Statement on Ethical Conduct in Research Involving Humans which builds on the ethical codes of the Declaration of Helsinki and the Principles of International Conference on Harmonisation Good Clinical Practice (as adopted in Australia). The study product and protocol were reviewed and approved by the Drug and Therapeutics Committee Clinical Trials Group (subcommittee of the WCHN Drug and Therapeutics Committee, approval date 17/10/2019). All study procedures and materials were reviewed and approved by the Women's and Children's Health Network Human Research Ethics Committee (approval No. HREC/19/WCHN/140, approval date: 22/11/2019) as well as the research governance officer at the Women's and Children's Hospital prior to commencement. Any change to the protocol, informed consent form or other study materials or procedures were considered amendments and were submitted to the governing ethics committee for approval prior to becoming effective. Any protocol deviations were noted in participant's records and were discussed by the steering committee.

The IMAGINE Trial is registered on the Australia and New Zealand Clinical Trial Registry (ANZCTR: ACTRN12620000552987) and the World Health Organisation International Clinical Trials Registry Platform (Universal Trial Number U1111-1250-1917).

Caregivers were provided with a participant information sheet about the IMAGINE trial and provided written informed consent for their infant's involvement prior to enrolment. Caregivers were free to renegotiate consent for each procedure in the study and could decline any part of the study, pause involvement in the study or withdraw their infants from the study at any time either verbally or in writing. If participants were non-compliant (in either the formula groups or the breastfeeding group), caregivers were encouraged and supported to remain in the study.

Caregivers were offered AUD\$40 (6 and 9 months) or AUD\$60 (12 and 24 months) to reimburse travel costs for attending clinical appointments. Families that relocated away from the study site were reimbursed for travelling from their residence to the clinic as per the concurrent Australian Government Tax Office cents per kilometre rate,⁴⁸ and in some circumstances, a night of accommodation or return flights were provided to allow attendance at the 12 and 24-month assessments.

This protocol and the results of this study will be published in peer-reviewed journals. Primary and key secondary outcomes related to neurodevelopment,

tolerance and safety will form the main results paper. Publication of information related to this study in formats including, but not limited to, conference abstracts, posters or presentations, seminars, journal articles, public reports, social media and internet postings. No participants will be identified in the dissemination of study results and data collected are treated with confidence. Participating families who have not withdrawn will receive a lay report of the study findings.

Access to data

Individual participant data, including data dictionaries, may be shared after deidentification on reasonable request. Proposals to access the data must be scientifically and methodologically sound and must be reviewed and approved by the IMAGINE trial Steering Committee, the Study Sponsor and the Women's and Children's Human Research Ethics Committee. To gain access, data requestors need to sign a data access agreement. Proposals should be directed to Chair of the Steering Committee, Jacqueline Gould through email (Jacqueline.gould@sahmri.com).

DISCUSSION

This protocol details a trial of MFGM-enriched formula compared with a standard infant formula without added MFGM for formula-fed infants to determine the effect on cognition. As is the gold standard for formula trials, we included an exclusively breastfed reference group that received no study formula even if breastfeeding ceased.

We carefully selected a comprehensive series of global and specialised assessments of early cognitive development. Our primary outcome is assessed with the Bayley Scales of Infant Development (now fourth edition),²⁷ a well-established test of early development and the most widely used assessment of early development internationally. Specialised assessments have more recently become of interest to nutrition researchers as they target specific abilities during infancy and toddlerhood that may be more sensitive to a nutrient intervention in a generally healthy, nutritionally replete population.^{49–51} Additional outcomes included health, growth and tolerance, as per guidelines set out by European Society for Paediatric Gastroenterology Hepatology and Nutrition for collecting core data in nutrition trials in infants.⁵² All assessments were conducted by trained research staff according to standard operating procedures. Study procedures were documented and published on a trial register in line with the recommendations for conducting and reporting clinical trials.^{53–59}

The sample size for this trial is large, based on the power calculation for the primary outcome and allowing for some loss to follow-up, although loss of contact with study families will still contribute to attrition bias. While the formula-fed groups were double-blinded to group allocation, the breastfed reference

group was not blinded to their group status or to the study aim of investigating the effect of formula with MFGM. At the time of designing the IMAGINE trial, face-to-face appointments were planned for enrolment, 4, 9, 12 and 24 months of age. However, prior to commencing, the COVID-19 became a worldwide pandemic and the Australian government mandated restrictions on non-essential face-to-face contact, particularly in the healthcare setting, from February 2020 for >7 months. Appointments were adapted to be conducted virtually, requiring adoption of e-consent, interviews via phone and postage of study formula to avoid the potential to spread the virus. Due to the need for remote data collection, infant growth measurements that were intended to be conducted at each appointment by trained study staff using calibrated equipment²⁵ were not possible. Instead, growth measurements conducted by healthcare professionals at routine check-ups (where available) were reported by caregivers and may not be performed as consistently as planned or may be missing if the infant has not attended a routine health check. Outcomes that necessitated a face-to-face appointment required face masks were occasionally missed (attention assessments) or postponed (Bayley-IV). Although the attention assessments were an optional outcome, the impact of the COVID-19 restrictions increased the amount of missing data for this assessment and reduced the available sample size. Overall, some adaptations were advantageous by allowing remote data collection and preferred by participants, while others may have contributed to a larger amount of missing data or wider age range at outcome assessments.²⁵ Importantly, IMAGINE commenced after the COVID-19 outbreak, so that study procedures were consistent throughout the study.

Australian guidelines recommended that infants are exclusively breastfed until 6 months, with ongoing breastfeeding to 12 months or beyond.^{26 60} Although breastfeeding initiation is high in Australia, exclusive breastfeeding and ongoing breastfeeding drops rapidly, so that by 6 months more than half of infants are fed formula, and by 12 months more than 80% are.⁶¹ It is important that breastmilk substitutes are nutritionally replete, and results of this trial will provide robust evidence for the effect of including MFGM in infant formula.

Author affiliations

¹SAHMRI Women and Kids, South Australian Health and Medical Research Institute, North Adelaide, South Australia, Australia

²School of Psychology, The University of Adelaide, Adelaide, South Australia, Australia

³School of Agriculture, Food and Wine, The University of Adelaide, Glen Osmond, South Australia, Australia

⁴SAHMRI Women and Kids, South Australian Health and Medical Research Institute Limited, Adelaide, South Australia, Australia

⁵School of Public Health, The University of Adelaide, Adelaide, South Australia, Australia

⁶Department of Psychology, The University of Kansas, Lawrence, Kansas, USA

⁷Neonatal Medicine, Women's and Children's Health Network, North Adelaide, South Australia, Australia

⁸Fonterra Research and Development Centre, Palmerston North, New Zealand

⁹Hamilton, Stockholm, Sweden

¹⁰Department of Dietetics & Nutrition, The University of Kansas, Lawrence, Kansas, USA

¹¹SAHMRI Women and Kids Theme, South Australian Health and Medical Research Institute Limited, Adelaide, South Australia, Australia

¹²Discipline of Paediatrics, The University of Adelaide, Adelaide, South Australia, Australia

Acknowledgements We would like to thank the families who are generously participating in the IMAGINE trial, the IMAGINE Trial Data Safety and Monitoring Committee (paediatrician Prof Cameron Grant, clinical researcher Lynley Drummond, and biostatistician Barbara Kuhn-Sherlock), as well as biostatistician Dr Thomas Sullivan, the IMAGINE Trial Steering Committee (Gould, Makrides, Yelland, McPhee), Investigative Team (Gould, Makrides, Yelland, McPhee, Gibson, Colombo, Roberts, Shaddy, Bednarz), and research staff, particularly co-ordinators Kimberly Hamlyn, Lauren Williams, Aya Gharrah, Eleanor Gardner and Beth Kean.

Contributors JFG, MM, AJM, JC, SG and RAG conceived the study and proposed the trial design. JFG, MM, JC, SG and RAG obtained funding. LNY and JB designed the analyses and performed the sample size calculations. JFG, MM, LNY, DJS and JC drafted the protocol. JFG, MM, AJM, JC, SG, LNY, JB, RMR, DJS and RAG contributed to the refinement of the study protocol manuscript and approved the final manuscript. JFG is the guarantor of the manuscript.

Funding The project was funded by the sponsor, Fonterra Co-Operative Group Limited, Auckland, New Zealand, and funding is paid to the South Australian Health and Medical Research Institute. MM is supported by an Australian National Health and Medical Research Council Principal Research Fellowship ID: GNT 2016756. Study products are provided by Fonterra Co-Operative Group Limited. The Australian National Health and Medical Research Council has no role in the study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication and has no authority over any of these activities.

Competing interests The project was funded by the sponsor, Fonterra Co-Operative Group Limited, Auckland, New Zealand, and funding was paid to the South Australian Health and Medical Research Institute. MM is supported by an Australian National Health and Medical Research Council Principal Research Fellowship ID: GNT 2016756. Study products were provided by Fonterra Co-Operative Group Limited. The Australian National Health and Medical Research Council has no role in the study design; collection, management, analysis and interpretation of data; writing of the report; and the decision to submit the report for publication and have no authority over any of these activities. Honoraria have been paid to Dr Gould's institution to support conference travel by NuMega Ingredients. No other competing interests are declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval All study procedures and materials were reviewed and approved by the Women's and Children's Health Network Human Research Ethics Committee (approval No. HREC/19/WCHN/140, approval date: 22/11/2019) as well as the research governance officer at the Women's and Children's Hospital prior to commencement.

Provenance and peer review Not commissioned; externally peer-reviewed.

Data availability statement Individual participant data, including data dictionaries, may be shared after de-identification upon reasonable request. Proposals to access the data must be scientifically and methodologically sound and must be reviewed and approved by the IMAGINE trial Steering Committee, the Study Sponsor, and the Women's and Children's Human Research Ethics Committee. To gain access, data requestors need to sign a data access agreement. Proposals should be directed to Chair of the Steering Committee, Jacqueline Gould through email (Jacqueline.gould@sahmri.com).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability

of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Jacqueline F Gould <http://orcid.org/0000-0003-2810-6870>

Lisa N Yelland <http://orcid.org/0000-0003-3803-8728>

REFERENCES

- Lönnerdal B. Bioactive proteins in human milk: mechanisms of action. *J Pediatr* 2010;156:S26–30.
- Horta B, CG V. *Long-Term Effects of Breastfeeding: A Systematic Review*. Geneva, Switzerland, 2013.
- Belfort MB, Knight E, Chandarana S, et al. Associations of maternal milk feeding with neurodevelopmental outcomes at 7 years of age in former preterm infants. *JAMA Netw Open* 2022;5.
- Kramer MS, Aboud F, Mironova E, et al. Breastfeeding and child cognitive development: new evidence from a large randomized trial. *Arch Gen Psychiatry* 2008;65:578–84.
- Anderson JW, Johnstone BM, Remley DT. Breast-feeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 1999;70:525–35.
- Larroque B, Ancel P-Y, Marret S, et al. Neurodevelopmental disabilities and special care of 5-year-old children born before 33 weeks of gestation (the EPIPAGE study): a longitudinal cohort study. *Lancet* 2008;371:813–20.
- Marlow N, Wolke D, Bracewell MA, et al. Neurologic and developmental disability at six years of age after extremely preterm birth. *N Engl J Med* 2005;352:9–19.
- Wang B. Sialic acid is an essential nutrient for brain development and cognition. *Annu Rev Nutr* 2009;29:177–222.
- Hernell O, Timby N, Domellöf M, et al. Clinical benefits of milk fat Globule membranes for infants and children. *J Pediatr* 2016;173 Suppl:S60–5.
- Timby N, Domellöf E, Hernell O, et al. Neurodevelopment, nutrition, and growth until 12 mo of age in infants fed a low-energy, low-protein formula supplemented with bovine milk fat globule membranes: a randomized controlled trial. *Am J Clin Nutr* 2014;99:860–8.
- Tanaka K, Hosozawa M, Kudo N, et al. The pilot study: Sphingomyelin-fortified milk has a positive association with the Neurobehavioural development of very low birth weight infants during infancy, randomized control trial. *Brain Dev* 2013;35:45–52.
- Li X, Peng Y, Li Z, et al. Feeding infants formula with probiotics or milk fat Globule membrane: a double-blind, randomized controlled trial. *Front Pediatr* 2019;7:347.
- Nieto-Ruiz A, García-Santos JA, Bermúdez MG, et al. Cortical visual evoked potentials and growth in infants fed with bioactive compounds-enriched infant formula: results from COGNIS randomized clinical trial. *Nutrients* 2019;11:2456.
- Li F, Wu SS, Berseth CL, et al. Improved neurodevelopmental outcomes associated with bovine milk fat Globule membrane and Lactoferrin in infant formula: a randomized. *J Pediatr* 2019;215:24–31.
- Xia Y, Jiang B, Zhou L, et al. Neurodevelopmental outcomes of healthy Chinese term infants fed infant formula enriched in bovine milk fat Globule membrane for 12 months - a randomized controlled trial. *Asia Pac J Clin Nutr* 2021;30:401–14.
- Colombo J, Harris CL, Wampler JL, et al. Improved neurodevelopmental outcomes at 5.5 years of age in children who received bovine milk fat Globule membrane and Lactoferrin in infant formula through 12 months: a randomized controlled trial. *J Pediatr* 2023;261:113483.
- Jiang B, Xia Y, Zhou L, et al. Safety and tolerance assessment of milk fat Globule membrane-enriched infant formulas in healthy term Chinese infants: a randomised multicenter controlled trial. *BMC Pediatr* 2022;22:465.
- Ambrożej D, Dumycz K, Dziechciarz P, et al. Milk fat Globule membrane supplementation in children: systematic review with meta-analysis. *Nutrients* 2021;13:714.
- Raza GS, Herzig KH, Leppälüoto J. Invited review: milk fat Globule membrane-A possible panacea for neurodevelopment, infections, cardiometabolic diseases, and frailty. *J Dairy Sci* 2021;104:7345–63.
- Timby N, Domellöf M, Lönnerdal B, et al. Supplementation of infant formula with bovine milk fat Globule membranes. *Adv Nutr* 2017;8:351–5.
- Arija V, Jardi C, Bedmar C, et al. Supplementation of infant formula and neurodevelopmental outcomes: a systematic review. *Curr Nutr Rep* 2022;11:283–300.
- Timby N, Adamsson M, Domellöf E, et al. Neurodevelopment and growth until 6.5 years of infants who consumed a low-energy, low-protein formula supplemented with bovine milk fat Globule membranes: a randomized controlled trial. *Am J Clin Nutr* 2021;113:586–92.
- World Health Organisation. WHO child growth standards: published by the centres for disease control and prevention. 2009.
- Gould JF, Yelland LN, Gibson RA, et al. Protocol for a multicentre prospective observational study of families with full-term infants on postnatal wards and in the community to capture feeding practices across the first year of life: the mother infant Lactation questionnaire (MILQ) study. *BMJ Open* 2022;12:e066355.
- Gould JF, Best K, Netting MJ, et al. New Methodologies for conducting maternal, infant, and child nutrition research in the era of COVID-19. *Nutrients* 2021;13:941.
- NHMRC. Infant feeding guidelines: information for health workers. In: Ageing D, ed. *Canberra National Health and Medical Research Council*. 2012.
- Bayley N, Aylward GP. *Bayley Scales of Infant and Toddler Development* 4th Edition: Australian and New Zealand Standardised Edition (Bayley-4 A & amp; NZ). Bloomington, M.N: NCS Pearson, 2019.
- Colombo J, Kannass KN, Shaddy DJ, et al. Maternal DHA and the development of attention in infancy and toddlerhood. *Child Dev* 2004;75:1254–67.
- Colombo J, Carlson SE, Cheatham CL, et al. Long-chain polyunsaturated fatty acid supplementation in infancy reduces heart rate and positively affects distribution of attention. *Pediatr Res* 2011;70:406–10.
- Colombo J, Gustafson KM, Gajewski BJ, et al. Prenatal DHA supplementation and infant attention. *Pediatr Res* 2016;80:656–62.
- Colombo J, Zavaleta N, Kannass KN, et al. Zinc supplementation sustained normative neurodevelopment in a randomized, controlled trial of Peruvian infants aged 6–18 months. *J Nutr* 2014;144:1298–305.
- Colombo J, Shaddy DJ, Richman WA, et al. The developmental course of habituation in infancy and preschool outcome. *Infancy* 2004;5:1–38.
- Colombo J, Richman WA. Infant timekeeping: attention and temporal estimation in 4-month-olds. *Psychol Sci* 2002;13:475–9.
- Colombo J, Richman WA, Shaddy DJ, et al. Heart rate-defined phases of attention, look duration, and infant performance in the paired-comparison paradigm. *Child Dev* 2001;72:1605–16.
- Blaga OM, Colombo J. Visual processing and infant ocular latencies in the overlap paradigm. *Dev Psychol* 2006;42:1069–76.
- Frick JE, Colombo J, Saxon TF. Individual and developmental differences in disengagement of fixation in early infancy. *Child Dev* 1999;70:537–48.
- Colombo J, Cheatham CL. The emergence and basis of endogenous attention in infancy and early childhood. *Adv Child Dev Behav* 2006;34:283–322.
- de Onis M, Onyango AW, Borghi E, et al. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007;85:660–7.
- Squires J, Twombly E, Bricker D, et al. *Ages & Stages Questionnaires® Third*. Baltimore, MD: Paul H. Brookes Publishing Co, 2009:256.
- Marchman VA, Fenson L, Thal DJ, et al. *MacArthur-Bates Communicative Development Inventories (CDI) User's Guide and Technical Manual Third*. Baltimore, MD: Brookes Publishing Co, 2023.
- Lewis SJ, Heaton KW. Stool form scale as a useful guide to intestinal transit time. *Scand J Gastroenterol* 1997;32:920–4.
- Frankenburg WK, Coons CE. Home screening questionnaire: its validity in assessing home environment. *J Pediatr* 1986;108:624–6.
- Walter T, De Andraca I, Chadud P, et al. Iron deficiency anemia: adverse effects on infant psychomotor development. *Pediatrics* 1989;84:7–17.
- Baghurst PA, McMichael AJ, Wigg NR, et al. Environmental exposure to lead and children's intelligence at the age of seven years. the port pirie cohort study. *N Engl J Med* 1992;327:1279–84.
- Makrides M, Gibson RA, McPhee AJ, et al. Effect of DHA supplementation during pregnancy on maternal depression and neurodevelopment of young children: a randomized controlled trial. *JAMA* 2010;304:1675.

- 46 Gamble C, Krishan A, Stocken D, *et al.* Guidelines for the content of statistical analysis plans in clinical trials. *JAMA* 2017;318:2337–43.
- 47 Sullivan TR, White IR, Salter AB, *et al.* Should multiple imputation be the method of choice for handling missing data in randomized trials *Stat Methods Med Res* 2018;27:2610–26.
- 48 Australian Tax Office. Deductions for motor vehicle expenses-cents per kilometre method Australian government. 2022. Available: <https://www.ato.gov.au/Business/Income-and-deductions-for-business/Deductions/Deductions-for-motor-vehicle-expenses/Cents-per-kilometre-method/2022>
- 49 Wainwright PE, Colombo J. Nutrition and the development of cognitive functions: interpretation of behavioral studies in animals and human infants. *Am J Clin Nutr* 2006;84:961–70.
- 50 Cheatham CL, Colombo J, Carlson SE. N-3 fatty acids and cognitive and visual acuity development: methodologic and conceptual considerations. *Am J Clin Nutr* 2006;83:1458S–1466S.
- 51 Georgieff MK. Nutrition and the developing brain: nutrient priorities and measurement. *Am J Clin Nutr* 2007;85:614S–620S.
- 52 Aggett P, Agostoni C, Axelsson I, *et al.* Core data for nutrition trials in infants: a discussion document--a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* 2003;36:338–42.
- 53 Calvert M, Kyte D, Mercieca-Bebber R, *et al.* Guidelines for inclusion of patient-reported outcomes in clinical trial protocols: the SPIRIT-PRO extension. *JAMA* 2018;319:483–94.
- 54 Chan A-W, Tetzlaff JM, Altman DG, *et al.* SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200–7.
- 55 Chan A-W, Tetzlaff JM, Gøtzsche PC, *et al.* SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ* 2013;346:e7586.
- 56 Begg C, Cho M, Eastwood S, *et al.* Improving the quality of reporting of randomized controlled trials. *JAMA* 1996;276:637–9.
- 57 Moher D, Hopewell S, Schulz KF, *et al.* CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *J Clin Epidemiol* 2010;63:e1–37.
- 58 Moher D, Schulz KF, Altman DG, *et al.* The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *Ann Intern Med* 2001;134:657–62.
- 59 Jarrold K, Helfer B, Eskander M, *et al.* Guidance for the conduct and reporting of clinical trials of breast milk substitutes. *JAMA Pediatr* 2020;174:874–81.
- 60 NHMRC. Eat for health; infant feeding guidelines: summary. In: *National Health and Medical Research Council DoHaA*. Canberra, Australia: National Health and Medical Research Council, 2013.
- 61 AIHW. Australian national infant feeding survey: indicator results. In: *Welfare AloHa*. Canberra AIHW, 2010.